

UNICEF WORLD SUMMIT FOR CHILDREN: END-DECADE INDICATORS

APPENDIX **F**

World Summit for Children, End-Decade Indicators: Armenia 2000

BASIC INDICATORS		
Childhood mortality	Under-five mortality rate (per 1,000 births)	39.0
	Infant mortality rate (per 1,000 births)	36.1
Maternal mortality	Maternal mortality ratio (deaths per 100,000 live births)	NA
Childhood malnutrition	Percent underweight (children under 5 years)	2.6
	Percent stunted (children under 5 years)	13.0
	Percent wasted (children under 5 years)	2.0
Clean water supply	Percent of household residents with safe water supply	91.8
Sanitary excreta disposal	Percent of household residents with latrine or toilet	99.9
Basic education	Percentage of children entering first grade of primary school who reach grade 5	99.3
	Percentage of primary-school age children currently attending primary school	94.5
	Proportion of 7 year olds entering primary school	68.1
SUPPORTING INDICATORS		
Family planning	Contraceptive prevalence (married women)	60.5
	Contraceptive prevalence (all women)	39.0
Safe motherhood	Percent of births with medical prenatal care ¹	92.4
	Percent of births with medical assistance at delivery ²	96.8
Low birth weight	Percent of births at low birth weight (below 2500 grams)	6.5
Micronutrient intake	Percent of households with iodised salt	83.6
	Percent of women who had night blindness while pregnant with last child ¹	1.4
Breastfeeding	Percent of infants less than 4 months of age exclusively breastfed	44.6
	Percent of infants 12-15 months still breastfeeding	28.8
	Percent of infants 20-23 months still breastfeeding	12.5
	Percent of infants 6-9 months receiving breast milk and complementary foods	50.8
Vaccinations	Percent of children 12-23 months receiving tuberculosis vaccine before 1st birthday	96.0
	Percent of children 12-23 months receiving DPT vaccine	95.1
	Percent of children 12-23 months receiving polio vaccine	97.6
	Percent of children 12-23 months receiving measles vaccine	78.8
Diarrhea treatment	Percent of children with diarrhea in preceding 2 weeks who received ORT ³	33.0
	Percent of children with diarrhea in preceding 2 weeks who received more fluids and continued eating somewhat less/the same/or more food	39.7
Acute respiratory infection	Percent of children with acute respiratory infection taken to a health facility	26.2
Childcare	Percent of children 0-14 years not living with both biological parents	10.1
	Percent of children 0-14 years who are orphans (either parent dead)	3.9
HIV/AIDS	Percent of women who had knowledge of 2 correct ways of avoiding HIV infection	43.9
	Percent of women who identified 3 misconceptions about HIV/AIDS	11.7
	Percent of women who correctly identified all 3 means of mother-to-child HIV transmission	55.1
	Percent of women expressing a discriminatory attitude towards people with HIV or AIDS	9.3
	Percent of women who stated knowledge of a place to be tested for HIV	44.4
	Percent of women who reported being tested for HIV	6.5
¹ Refers to last birth in the five years preceding the survey ² Refers to all births in the five years preceding the survey ³ Includes ORS and/or increased fluids na = Not applicable		